



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The surface $OA'A$ is the total conical surface less the ungula, that is,

$$\frac{\pi}{2} \sin \alpha [2K'^2 - 2K^2 + (K + K') \sqrt{KK'}] = \frac{\pi}{2} (K + K')(KK')^{\frac{1}{2}} \sin \alpha.$$

The problem rests upon the tedious integration of the form,

$$\int_r^R 2x \cos^{-1} \left[\frac{2Rr - (R + r)x}{(R - r)x} \right] dx.$$

Also solved by A. M. HARDING and P. PENALVER.

A solution of 334 was received from ELMER SCHUYLER, after the the forms were sent to the printer.

MECHANICS.

270. Proposed by W. J. GREENSTREET, Burghfield, England.

A cycloid has its base vertical. Find the line of quickest descent from the middle point of the base, and its approximate inclination to the horizon.

SOLUTION BY J. SCHEFFER, Hagerstown, Md.

Let AB be any straight line drawn from the middle point A of base of the cycloid, and let θ be the angle that it makes with the horizon. The time of descent

from A to B is $t = \sqrt{\frac{2}{g}} \cdot \frac{s}{\sin \theta}$, AB being s . But $s^2 = (r\pi - x)^2 + y^2$, x and y

being subject to the equations of the cycloid, $x = r(\phi - \sin \phi)$, $y = r(1 - \cos \phi)$.

Substituting, we have $s^2 = r^2[(\pi - \phi) + \sin \phi]^2 + (1 - \cos \phi)^2$.

We have to get the minimum of $s/\sin \theta$, which reduces to

$$\frac{(\psi + \sin \psi)^2 + (1 + \cos \psi)^2}{\psi + \sin \psi},$$

after putting $\pi - \phi = \psi$ and omitting the constant r^2 .

Differentiating and setting the differential coefficient equal to zero, we get, after some easy reductions, the transcendental equation $\psi^2 - 2 \cos \psi - 2 = 0$. Whence $\psi = \pm 2 \cos \psi/2$, and using the positive sign, we find $\psi = 85^\circ$, nearly. Then

$$\tan \theta = \frac{r\pi - x}{y} = \frac{\psi + \sin \psi}{1 + \cos \psi}.$$

Hence, finally, $\theta = 53^\circ 39'$, nearly.

Note. It is assumed in the above solution that the line sought is the *straight line* of quickest descent. Otherwise the problem becomes much more difficult, requiring for its solution the calculus of variations. THE EDITORS.

NUMBER THEORY.

A solution of 188 was received from ELMER SCHUYLER after the December issue had been made up.

